


# Micronutrient mixture formulated in water soluble powder

# Combi-Mix



## Net Weight

1 Kg

5 Kg

25 Kg

## Contains

<b>Iron</b>	<b>4.10%</b>
<b>Magnesium</b>	<b>4.60%</b>
<b>Manganese</b>	<b>3.80%</b>
<b>Zinc</b>	<b>1.40%</b>
<b>Boron</b>	<b>0.65%</b>
<b>Copper</b>	<b>1.30%</b>
<b>Molybdenum</b>	<b>0.07%</b>
<b>Sulphur</b>	<b>4.40%</b>

E.C. Fertilizer

**Combi-Mix** is a micronutrient mixture for the prevention and some times the correction of deficiencies in many plants. It contains 4.10% Iron (Fe), 1.40% Zinc (Zn), 3.80% Manganese (Mn), 4.60% Magnesium (Mg), 0.65% Boron (B), 1.30% Copper (Cu), 0.07% Molybdenum (Mo) and 4.40% Sulphur, all derived from complex chelated compounds (chelating agent EDTA) and mineral salts. **Combi-Mix** is totally water soluble and is recommended mainly for foliar applications in all crops. **Combi-Mix** is essential in order to maintain the health and the high yields for all intensive crops.

## APPLICATIONS - APPLICATION PERIOD

**Foliar application:** Foliar application is recommended for the quick correction of deficiencies. In order to achieve maximum benefits, foliar applications should start at the beginning of the growing season when there is sufficient foliage to absorb the spray. In case of severe deficiencies, applications must be repeated every 7-14 days. When applied foliarly on its own, it is recommended to add a sticking agent in the solution.

**Soil application:** Dissolve in an appropriate amount of water, then apply as a course low pressure spray, or through drip irrigation along with a liquid fertilizer. Apply in the soil close to plants and tree trunks. Work the product well into the soil after application and water carefully after each application as to enhance nutrient uptake.

## COMPATIBILITY

**Combi-Mix** can be mixed with most fertilizers and agrochemicals without creating sediment or causing scorch problems. When mixed with liquid fertilizers the mixture should be applied as soon as possible.

## PRECAUTIONS

Keep well sealed and store in its original sealed container. Wash thoroughly your hands before meals and after handling. Keep out of reach of children and pets. Store dry and away from strong light. Do NOT apply to crops suffering stress such as extreme temperatures, strong winds etc. Avoid swallowing and contact with eyes and skin. In case of eye irritation, flush with plenty of water.

PRODUCED BY

**HUMOFERT**



1 Ermou & Theotokopoulou, Metamorphosis, Greece

Tel. +30 210 284 5891 Fax. +30 210 281 7971

E-mail: info@humofert.gr Web Site: www.humofert.gr

## APPLICATION RATES

The following application rates are general. Precise application rates must be determined according to the type and the size of the plant.

### Foliar application

Open field vegetables : 0.5 - 1 kg/ha.

Cereals : 1 - 2 kg/ha.

Vineyards : 1 - 1.5 kg/ha.

Fruit trees (apples, pears etc.) : 1 - 2.5 kg/ha. It must not be applied on apples of Golden variety.

Bushes (strawberries, blackberries etc.) : 1 - 1.5 kg/ha.

Greenhouse vegetables (tomatoes, cucumbers, peppers etc.) : 0.5-1 kg/ha.

Ornamentals : 0.5 - 1 kg/ha.

**Note :** Concentration must never exceed 0.1%w/v. It must not be applied at full bloom.

### Soil application

Vegetables : 4 - 8 kg/ha.

Cereals : 4 - 10 kg/ha.

Fruit trees & bushes : 4 - 10 kg/ha.

Ornamentals : 2 - 10 kg/ha.

## DANGER



**Hazard Statements:** H360FD: May damage fertility. May damage the unborn child.

**Precautionary statements:** P201: Obtain special instructions before use. P281: Use personal protective equipment as required. P308+P313: IF exposed or concerned: Get medical advice/attention. P405: Store locked up. P501: Dispose of contents/container to ... in accordance with local/regional/national/international regulation (to be specified).

**Contains:** Disodium Octaborate Tetrahydrate

**FOR PROFESSIONAL USE ONLY**