


Natural Condensed Biostimulant

Perfect

For vigorous growth, increased yields
and improved quality of harvest
in all crops



Produced by



1 Ermou & Theotokopoulou str., Metamorphosis, Greece
Tel. 210 284 5891 - Fax. 210 281 7971
E-mail: info@humofert.gr Web Site: www.humofert.gr

Net Content

1 L

APPLICATION RATES - APPLICATIONS

Vegetables - Horticulture - Sweet corn

Foliar spray : 4 - 5 l / ha. **Fertigation :** 25 - 30 l / ha.

Applications: 1) At the emergence of the first leaves (4-6 leaves) or for sweet corn when plants reach 20 - 25 cm height. 2) Before blooming or prior to tasseling for sweet corn. 3) At the first fruit set. 4) Before harvest. 5) 48 hours after each picking.

Cereals - Cotton - Tobacco - Beans - Potato

Foliar spray : 2.5 - 5 l / ha. **Fertigation :** 10 - 20 l / ha.

Applications: 1) At the emergence of the first leaves for Cotton-Tobacco-Beans-Potato, when plants reach 10-13 cm height for cereals or when plants reach 20 - 25 cm height for corn. 2) Before blooming or prior to tasseling. 3) Before harvest.

Note: For potatoes it is recommended to dip the potato seeds in a solution 1 l / 100 l of water for 24 hours before planting.

Tree crops

Foliar spray : 5 - 7 l / ha. **Fertigation :** 30 - 50 l / ha.

Applications: 1) Prior to blooming or at the beginning of blooming. 2) At petal fall (after the fall of the 2/3 of the petals). 3) At fruit set. 4) Every 15 days until harvest. 5) After harvest.

Vineyard

Foliar spray : 4 - 8 l / ha. **Fertigation :** 30 - 40 l / ha.

Applications: 1) Prior to blooming or at the beginning of blooming. 2) At berry set, during shattering. 3) At fruit sizing (2nd & 3rd sizing stage). 4) Before harvest.

Alfalfa-Clover-Fodders

Foliar spray : 4 l / ha. **Fertigation :** 10 - 20 l / ha.

Applications: 1) Early in the spring. 2) After each cutting or pasturing.

Ornamental plants

Foliar spray : 1 l / 250 l of water. **Fertigation :** 25 - 30 l / ha.

Applications: Every 2 - 3 weeks.

USE

Shake well before each use. Apply within 24 hours after the product's dilution. Apply directly in the tank.

COMPATIBILITY

It can be mixed with most herbicides, insecticides, fungicides and foliar fertilizers. In any case a test of compatibility is recommended before mixing in the tank.

Approved for use in organic agriculture according to Reg. (EC) 889/2008.

KEEP OUT OF REACH OF CHILDREN

